

Infrared Spectroscopy of U Equulei's Warm Circumstellar Gas

T. R. Geballe¹, C. Barnbaum², Keith S. Noll³, and M. Morris⁴

Received _____; accepted _____

to be published in the Astrophysical Journal, Vol. 625, May 20, 2005

¹Gemini Observatory, 670 N. A'ohoku Place, Hilo, HI 96720; tgeballe@gemini.edu

²Department of Physics, Astronomy, & Geosciences, Valdosta State University; Valdosta, GA 31698; cbarnbaum@valdosta.edu

³Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21218; noll@stsci.edu

⁴Department of Physics and Astronomy, University of California at Los Angeles, Los Angeles, CA 90095; morris@astro.ucla.edu

ABSTRACT

Medium and high resolution spectroscopy of U Equulei from 1 to 4 μm during 1997-2003 has revealed information about its unusual circumstellar envelope, observed previously at optical and radio wavelengths. Strong absorption bands of H_2O and of CO dominate the 1–4 μm spectrum. The gas has a mean temperature of 600 K and $^{12}\text{C}/^{13}\text{C} \leq 10$. The CO 2-0 line profiles and velocities imply no net ejection or infall and indicate either rapid radial gas motions being seen along a narrow continuum beam, or absorption by orbiting gas that is nearly coincident with a highly extended continuum source. The gas could be located in a disk-like structure. The observed high column densities of warm CO and H_2 normally would be associated with sufficient dust to completely obscure the star at optical wavelengths. The observations thus indicate either a highly abnormal gas-to-dust ratio, consistent with the earlier optical observation of abundant refractory metal oxides in the circumstellar gas, or peculiar geometry and/or illumination.

Subject headings: stars: U Equ; circumstellar matter - infrared: stars - molecular processes

1. Introduction

The peculiar variable star, U Equulei (IRAS 20547+0247), has been described by Barnbaum et al. (1996) as having one of the most unusual optical spectra ever observed. Their spectra, obtained in 1994, are dominated by strong absorption bands of metallic oxides, superficially similar to those of mid-late M giants. However the TiO, AlO and VO in U Equ are simultaneously in absorption and in emission. In addition, the bands of these molecules are finely structured, implying a circumstellar, rather than photospheric origin for them. The absorption features must arise from a region so cool that the higher rotational levels are not populated. Yet, at such a temperature the titanium seen in TiO should be in grains.

Besides its unique optical spectrum U Equ displays other peculiarities. Its OH and H₂O maser emission (Sivagnanam et al. 1990 and Zuckerman & Lo 1987, respectively) vary in velocity as well in strength. U Equ has a large radial velocity of -78 km s^{-1} (LSR, -91 km s^{-1} heliocentric; Sivagnanam et al. 1990); the uncertainty in this value may be as much as $10\text{--}20 \text{ km s}^{-1}$ (C. Barnbaum 1996, unpublished data). The object is also at high galactic latitude (-26°), suggesting a possible link to a halo population. The $25/12 \text{ }\mu\text{m}$ color observed by the Infrared Astronomical Satellite (IRAS) is typical of optically thin circumstellar envelopes, despite its IRAS LRS spectrum displaying a moderately strong $10 \text{ }\mu\text{m}$ silicate absorption band (Barnbaum et al. 1996, Chen & Gao 2002), indicating a dusty envelope. Almost all other oxygen-rich stars with similar $25/12 \text{ }\mu\text{m}$ colors show the $10 \text{ }\mu\text{m}$ feature in emission, consistent with optically thin shells.

Barnbaum et al. (1996) suggested that U Equ’s silicate absorption feature arises in a thick, dusty circumstellar disk. Since the $25/12 \text{ }\mu\text{m}$ colors indicate optically thin dust, they proposed that the dusty disk is seen edge-on, with an extended spherical or asymmetrical component giving rise to the 12 and $25 \text{ }\mu\text{m}$ emission.

U Equ is listed in the General Catalogue of Variable Stars as an irregular long period variable (Lb) with a magnitude variation in the P (photographic) band of 14.5 to 15.5. Barnbaum et al. (1996) estimate its distance as 1.5 kpc. In 1994, the visual magnitude was approximately 9, yet when it was observed again in 1996, 1998, and 1999, the star had faded to $V=13$. The variations in the the visual magnitude and the origin of the circumstellar material are unexplained. However, the unusual nature of the optical spectrum suggests that U Equ might have entered a rapid stage of evolution. Siess & Livio (1999) have considered that accretion of a giant planet onto a growing red giant could spin up the stellar envelope sufficiently to extrude an equatorially compressed outflow, or expanding disk, and that this might resemble the peculiar environment of U Equ.

In this paper we report infrared spectroscopy of U Equ, which allows a fuller characterization of the physical conditions in the circumstellar material. In particular and in corroboration of the optical study of Barnbaum et al. (1996), we also have found a large quantity of absorbing gaseous matter that is difficult to reconcile with the relatively modest amount of extinction by dust.

2. Observations

Spectra of U Equ in the 1.0–2.5 and 2.9–4.1 μm regions were obtained at the 3.8 m diameter United Kingdom Infrared Telescope (UKIRT) at various times during 1997–2003. An observing log is provided in Table 1. The medium resolution observations employed UKIRT’s facility spectrographs: CGS4 (Mountain et al. 1990) with its 40 and 150 l/mm gratings, and 0.6'' wide slit; and UIST (Ramsay Howat et al. 2000) with its HK grism and 0.48'' wide slit. Resolving powers ranged from 450 to 4100. The spectra were obtained in the stare - nod-along-slit mode. The calibration stars listed in the table were observed immediately before or after U Equ for the purpose of flux calibration and removal of telluric

absorption lines. Fluxes of stars in the various photometric bands were estimated from their visual magnitudes and visible-infrared colors predicted for their spectral types (Tokunaga 2000). Wavelength calibration was derived from spectra of arc lamps and is accurate to considerably better than $0.001 \mu\text{m}$. The echelle in CGS4 was also used in 2003 to obtain high resolution spectra in a number of narrow wavelength intervals in the K window at resolving powers of about 20,000, corresponding to a velocity resolution of 15 km s^{-1}). For these spectra wavelength calibrations were obtained from telluric absorption lines and are accurate to 3 km s^{-1} .

Data reduction employed Figaro routines to extract source spectra from the spectral images produced by the CGS4 and UIST arrays, wavelength-calibrate them, ratio them, and perform an initial flux calibration. It was found that where the lower resolution *JHK* and *LL'* spectral segments overlapped or adjoined one another flux levels disagreed typically by 20 percent. The mismatches are probably due to variations in guiding accuracy and in the seeing (which usually was comparable to the slit width) during the observations, as well as to inaccuracies in the IR magnitudes of the calibration stars used in the reduction. Another possibility is variations of U Equ itself. However, the two 3-4 μm segments, which were obtained more than two years apart, were mismatched by only 20%. The spectral segments were scaled by small factors in order to adjoin them smoothly. The final flux calibrations are believed to be accurate to $\pm 20\%$.

3. Results

3.1. Overview of 1-4 μm Spectrum

Figure 1 shows the medium resolution 1.0–2.5 μm spectrum of U Equ as observed in 1997 together with the 2.9–4.1 μm spectrum as observed in 1998 and 2000. The

1.4–2.5 μm spectrum obtained in 2003 is closely similar, in depths of absorption bands and in overall continuum slope, to the spectrum in the top panel of Fig. 1. The 1-4 μm spectrum is dominated by water vapor absorption, which is responsible for the deep bands at 1.1–1.2 μm , 1.3–1.5 μm and 1.8–2.0 μm as well as the detailed spectral structure at 2.9–3.4 μm . Weak spectral structure is also evident longward of 3.7 μm (see Fig. 1, lower panel), but we are unable to identify the absorbing species at this spectral resolution and signal-to-noise ratio (~ 50). Also apparent in the spectrum is absorption due to the first overtone band of carbon monoxide at 2.3–2.4 μm . The 1.8–2.0 μm water band had been seen previously in a 1.6-2.5 μm survey spectrum by Lancon & Wood (2000), but the CO absorption was not reported by those authors, although in retrospect it appears to be present in their spectrum at a low signal-to-noise ratio.

The shape of the 2.3–2.4 μm spectrum in Fig. 1, with its local maximum at the CO 2-0 band center near 2.343 μm , indicates that most of the CO that is in the lowest ($v=0$) vibrational state. Thus, the CO is much cooler than a stellar photosphere and must reside in circumstellar material. The narrowness of the 1.9 μm water band in comparison to those of cool Miras (e.g., see Lancon & Wood 2000) also suggests that the absorbing material is quite cool.

3.2. Higher resolution spectra: CO, H₂O, and H₂

A higher resolution ($R \sim 4100$) spectrum of the CO first overtone band obtained in 1998 is shown in Fig. 2. In it the vibration-rotation structure of the P and R branches 2-0 band of CO is clearly resolved into lines. The strongest lines correspond to $J \sim 10$, but individual lines in the R branch can be seen out to $J=25$ at this resolution. As in the medium resolution spectrum, no 2-0 band head (corresponding to $J \sim 50$) is apparent at 2.293 μm . The unevenness of the band structure indicates that absorbers and possibly

emitters other than CO are present as well, especially near 2.320–2.325 and 2.340–2.345 μm . ^{13}CO and other rarer isotopic species of CO cannot be significant contributors in either of these spectral intervals.

Fig. 3 shows the portion of the R=4100 spectrum from 1998 in more detail, as well as the same interval observed with the CGS4 at R=18000 in 2003. This region contains the strongest CO lines. The difference in the observed widths and depths of the lines is consistent with the different resolutions. From the 2003 spectrum we derive a deconvolved FWHM of 35 km s $^{-1}$ for these lines. At both times the lines were centered at the nominal radial velocity, -91 km s $^{-1}$ hel. At the higher resolution of the 2003 spectrum several weaker lines are present. With the aid of the HITRAN2004 database (Rothman et al. 2003) we have identified them as H $_2$ O from relatively low-lying energy levels.

Fig. 4 is an echelle spectrum in the vicinity of the CO 2-0 band head. As was apparent in the lower resolution spectra of Figs. 1 and 2, the band head is not seen. The highest J levels that can be distinguished are near J=30. H $_2$ O accounts for most of the strong absorption lines in this spectral interval.

The spectrum in Fig. 5 covers an interval near 2.38 μm containing the strongest lines of the 2-0 R branch of ^{13}CO . Most of the ^{13}CO lines are blended with stronger lines of H $_2$ O and/or the ^{12}CO 2-0 P branch and are not readily apparent. However, the R(10) and R(11) lines of ^{13}CO are fairly well isolated and clearly detected and the R(12) line also is apparent between lines of the other species. Equivalent widths cannot be determined with accuracy because of uncertainties both in the amounts of line blending and in the placement of the continuum.

A high resolution spectrum also was obtained at the wavelength of the 1-0 S(1) transition of H $_2$ and is shown in Fig. 6. This line was detected and its equivalent width over a 75 km s $^{-1}$ interval centered at the stellar velocity measured to be $9 \pm 2 \times 10^{-6}$ μm . The

core of the line appears to be noticeably narrower than the CO, but this could easily be a consequence of the low signal-to-noise ratio of the spectrum.

4. Discussion

4.1. CO Temperature, $^{12}\text{C}/^{13}\text{C}$, and Column Densities

We have attempted to fit the ^{12}CO absorption spectrum in Fig. 2 with a multilayer slab models. Spectra of several single temperature slabs are shown in Fig. 7. Even though other absorbing species, principally H_2O , contaminate the spectrum it is clear that at temperatures between 500 K and 1000 K this simple model gives a good general representation of the CO absorption. If the central star is a mid-G giant as suggested by Barnbaum et al. (1996), and all of the CO is at 600 K, the CO is located roughly 5×10^{13} cm (3 a.u.) from the star. At a distance of 1.5 kpc, a disk or shell of this dimension subtends only 0.005'' on the sky and would not be resolved by adaptive optics on the largest telescopes. Indeed upper limits to the diameter from HST imaging (C. Barnbaum 1996, unpublished data) and Keck adaptive optics imaging (M. Morris 2004, unpublished data) are each $\sim 0.05''$.

Figure 8 shows four-layer (2000K, 1500K, 1000K, and 500K) model spectra near $2.38 \mu\text{m}$, the same interval shown in Fig. 5 containing isolated lines of ^{13}CO and numerous lines of H_2O . Each synthetic spectrum has a ^{12}CO column density of $2 \times 10^{20} \text{ cm}^{-2}$, which fits the ^{12}CO line strengths well. The upper panel shows the effect of varying the H_2O column density for $^{12}\text{C}/^{13}\text{C} = 4$, whereas the lower panel shows the effect of varying $^{12}\text{C}/^{13}\text{C}$. As noted previously many of the CO lines are blended with H_2O lines; in addition either the H_2O line list is incomplete or there are other contaminants. Nevertheless, adding H_2O noticeably improves the fit, and these simple models suggest that $^{12}\text{C}/^{13}\text{C} \sim 4$ and

that H₂O is approximately 2.5 times more abundant than CO.

For the above CO column density and temperature, and at the observed line width, the strongest lines of ¹²CO have optical depths at line center of order unity in the lowest temperature layer, which contains roughly half of the CO. Therefore the best fit model spectrum with ¹²C/¹³C = 4 probably only moderately underestimates ¹²C/¹³C, and we conclude that the ratio is ≤ 10 . Assuming U Equ has $[C]/[H] = 2.6 \times 10^{-4}$ (Asplund 2003) and that all carbon is in CO, the measured CO column density implies $N(H_2) \sim 4 \times 10^{23} \text{ cm}^{-2}$. Using the formulae in Lacy et al. (1994) the measured equivalent width of the H₂ S(1) line yields an H₂ column density of $4 \times 10^{23} \text{ cm}^{-2}$ in the $v=0, J=1$ state, which is roughly half the total column density of H₂ at temperatures of 500–1000 K. The comparison suggests that gas phase carbon might be depleted by a factor of roughly 2.

The column densities of CO and H₂ in U Equ’s circumstellar material are remarkably high. Justannont et al. (1996) found a CO column density of $1 \times 10^{19} \text{ cm}^{-2}$ in the circumstellar gas of the oxygen-rich mass-losing supergiant NML Cygni and Hall & Ridgway (1978) found $1 \times 10^{19} \text{ cm}^{-2}$ of CO in the mass-losing carbon star IRC+10216. The optical extinctions to these two evolved stars are ~ 15 and ~ 20 magnitudes, respectively (Ridgway et al. 1986; Ivesić & Elitzur 1996), much greater than the 2.1 mag toward U Equ (Barnbaum et al. 1996). Column densities of warm CO observed in the dense winds of young stellar objects (e.g., Mitchell et al. 1990) typically are more than two orders of magnitude less than toward U Equ. In interstellar material a gas column density such as that observed in U Equ would be associated with a visual extinction of several tens of magnitudes as well as an H₂ column density of $\sim 10^{23} \text{ cm}^{-2}$ (using the cosmic value of $[C]/[H]$). The former is in conflict with the estimate of Barnbaum et al. (1996) and is clearly unrealistic in view of the visual brightness of U Equ.

Thus, it appears that either the gas-to-dust ratio is much higher in U Equ than in

typical circumstellar or interstellar matter, or the dust and associated gas are located in a thin disk that transmits or scatters (above and below the disk) a significant fraction of the photospheric light. In the latter case the above column density is a lower limit.

4.2. Velocities

The CO line profiles, which are centered on the stellar velocity and are symmetric about it, show no evidence for net infall or outflow of the circumstellar material. This result is consistent with the OH line measurements by Sivagnaman et al. (1990), which indicate radial motions of only a few km s^{-1} . A gaseous shell at $T \sim 600$ K stably orbiting U Equ at radius $\sim 5 \times 10^{13}$ cm, and observed in absorption against a central star, would produce much narrower absorption lines than the observed 35 km s^{-1} FWHM. If the gas is in a shell-like structure, its motion is highly turbulent and has maintained that turbulence over 6 years, which does not seem likely.

The orbital speed of material at the inferred distance of $\sim 5 \times 10^{13}$ cm from a roughly solar mass star is 20 km s^{-1} . Thus, the observed 35 km s^{-1} linewidths suggest that the infrared continuum could be highly extended and nearly coincident with the absorbing gas. The continuum would then result from scattering, absorption, and reemission of photospheric radiation. This crude model probably can account for the observed broad lines but cannot explain the low extinction.

5. Conclusion

Like the optical spectrum, the infrared spectrum of U Equ appears to be unique. Its most remarkable qualities are the presence of large amounts of warm circumstellar CO and water vapor and, as surmised from the optical spectrum, an extreme lack of obscuring dust.

The abundances in U Equ are not well quantified, but certainly the presence in the optical spectrum of strong TiO and other metallic oxide features suggests that the ingredients for forming copious amounts of dust are present. The temperature of the CO is well below the dust condensation temperature of metal oxides. Thus all evidence to date suggests that U Equ should be surrounded by a totally obscuring cloud of dust. That this is not the case probably either implies a remarkably high gas to dust ratio or peculiar geometry and illumination, perhaps with starlight illuminating and heating the edge of a very thin circumstellar disk-like structure.

It has been proposed that the phenomena being seen in U Equ might be consistent with the destruction of a planet. Siess & Livio (1999) suggested that U Equ might be extruding metal- enriched material because of rapid rotation caused by the accretion of a giant planet. However, there is no evidence for ejection (i.e., no net outflow of gas; indeed not even asymmetry in the line profiles). Moreover, the very low value of $^{12}\text{C}/^{13}\text{C}$ in the circumstellar material indicates that material is not highly diluted by planetary matter, which would have a high value of $^{12}\text{C}/^{13}\text{C}$. Mass estimates are very uncertain because of the unknown geometry of the absorbing CO. If $[\text{C}]/[\text{H}]$ is solar and even if the observed CO is in a spherical shell of radius 5×10^{13} cm, the mass of the shell is only 0.03 that of Jupiter, which, when combined with the observed value of $^{12}\text{C}/^{13}\text{C}$ again suggests that there has been very little dilution by a planet.

Although the current study provides information on the evolutionary state and some of the general properties of its circumstellar material, it has not solved the enigma that is U Equ. In particular, the tools of very high resolution spectra, polarimetry, and interferometry should now be brought to bear on this most fascinating object.

We wish to thank the staff of the Joint Astronomy Centre for its support of these observations. UKIRT is operated by the Joint Astronomy Centre on behalf of the U.K.

Particle Physics and Astronomy Research Council. TRG’s research is supported by the Gemini Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., on behalf of the international Gemini partnership of Argentina, Australia, Brazil, Canada, Chile, the United Kingdom and the United States of America.

REFERENCES

- Asplund, M., 2003 in CNO in the Universe, A.S.P. Conf Ser. 304, ed. C. Charbonnel, D. Schaerer & G. Meynet (San Francisco: ASP), 10
- Barnbaum, C., Omont, A. & Morris, M. 1996, A&A, 310, 259
- Chen, P.-S. & Gao, Y.-F. 2002, Chin. J. Astron. Ap., 2, 169
- Hall, D. N. B. & Ridgway, S. T. 1978, Nature, 273, 281
- Ivesić, Z. & Elitzur, M. 1996, MNRAS, 279, 1019
- Kholopov, P. N. 1985, General Catalog of Variable Stars (4th Edition, Moscow: Nauka)
- Justannont, K., et al. 1996, A&A, 315, L217
- Lacy, J. H., Knacke, R., Geballe, T. R. & Tokunaga, A. T. 1994, ApJ, 428, L69.
- Lancon, A. & Wood P. R. 2000, A & A Suppl., 146, 217
- Mitchell, G. F., Maillard, J.-P., Allen, M., Beer, R. & Belcourt, K. 1990, ApJ, 363, 554
- Mountain C. M., Robertson D. J., Lee T. J. et al. 1990, SPIE 1235, 25
- Ramsay Howat, S. K., Ellis, M. A., Gostick, D. C., Hastings, P. R., Strachan, M., & Wells, M. 2000, Proc. SPIE, 4008, 1067
- Ridgway, S. T., Joyce, R. R., Connors, D., Pipher, J. L. & Dainty. C. 1986, ApJ, 302, 662
- Rothman, L. S., et al. 2003, J. Quant. Sp. Rad. Trans., 82, 5
- Siess, L. & Livio, M. 1999, MNRAS, 308, 1133
- Sivagnanam, P., le Squeren, A. M., Minh, F. Tran & Brax, M. A. 1990 A&A, 233, 112
- Tokunaga, A. T., 2000, in "Allen's Astrophysical Quantities," ed. A. N. Cox, (New York: Springer Verlag), 143

Zuckerman, B. & Lo, K. Y. 1987, A&A, 173, 263




Fig. 1.— Medium resolution 1–4 μm spectrum of U Equ. Prominent absorption bands are identified.




Fig. 2.— Spectrum of U Equ in the 2–0 band of CO. The P and R branches, band center, and the unseen band head at $2.2927 \mu\text{m}$ are indicated.




Fig. 3.— Details of the CO 2-0 R branch lines as observed in 1998 and 2003. The wavelength scale is for zero heliocentric velocity. Lines of CO and known strong lines of H₂O are identified; their positions, indicated by vertical lines, are shifted by -91 km s^{-1} , the heliocentric velocity of U Equ. Velocity resolutions are also shown.




Fig. 4.— High resolution spectrum near the CO 2-0 band head. Positions of identified lines of CO and H₂O are shifted by -91 km s^{-1} . The location of the band head is also shown.




Fig. 5.— High resolution spectrum in the ^{12}CO 2-0 P branch, showing detection of R(10) - R(13) lines of ^{13}CO . All line indicators are shifted by -91 km s^{-1} ; known strong water lines are indicated by dashed lines.




Fig. 6.— Spectrum at the wavelength of the H₂ 1-0 S(1) line. The noise level, spectral resolution, and line wavelength, Doppler-shifted by -91 km s⁻¹ are indicated.




Fig. 7.— The observed R=4000 spectrum of U Equ in the CO 2-0 band (top) compared with isothermal slab absorption models of a CO column density of $1 \times 10^{20} \text{ cm}^{-2}$ at four temperatures.




Fig. 8.— The observed R=19000 spectrum of U Equ near $2.38 \mu\text{m}$ (thick line) compared with multislabs synthetic spectra with $N(^{12}\text{CO}) = 2 \times 10^{20} \text{ cm}^{-2}$ and various amounts of ^{13}CO and H_2). Most of the strong lines are due to ^{12}CO . Relatively unblended lines of ^{13}CO are indicated. The upper panel shows spectra with $^{12}\text{CO}/^{13}\text{C} = 4$ and $[\text{H}_2\text{O}]/[\text{CO}] = 0$ (dotted line) and 2.5 (thin line). The lower panel shows spectra with $^{12}\text{C}/^{13}\text{C} = 8$ (dotted)

Table 1. Observing Log

Date	Wavelength (μm)	R	Exposure (s)	Calib. star
19970802	1.00-1.33	950	288	HR 7973 (F5V)
19970802	1.44-2.10	700	96	HR 7973 (F5V)
19970802	1.87-2.52	880	96	HR 7973 (F5V)
19980407	2.25-2.42	2100	1080	HR 7973 (F5V)
19980422	2.29-2.37	4100	1080	HR 8205 (A1V)
19980624	3.45-4.09	1500	240	HR 8328 (A1V)
20001208	2.89-3.52	1280	576	HR 8205 (A1V)
20030824	1.40-2.50	450	1800	HIP 102805 (F5V)
20030824	2.328-2.344	18000	960	HR 8178 (A3V)
20030824	2.115-2.127	21000	1920	HIP 102631 (A0V)
20030909	2.374-2.388	19000	480	HIP 102631 (A0V)
20031113	2.292-2.305	21000	1440	HIP 102631 (A0V)